

# The Mediating Role of Duration and Intensity of Physical Activity in Increasing Eating Awareness: Adolescent and Parental Participation Study

Mustafa AKIL

Faculty of Sport Sciences, University of Usak, Turkey

Corresponding author: mustafa.akil@usak.edu.tr

**Abstract** Mindful eating is a condition in which a person becomes aware of the moment while eating. Very few studies have investigated eating awareness in environments where physical activity is performed. The aim of this study is to investigate whether the duration and intensity of physical activity in adolescents aged 15–17 affects their nutritional awareness. 1117 adolescents aged 15–17 and one parent of each were involved in the study. It is a cross-sectional study. Personal information form, International Physical Activity Questionnaire Short Form (IPAQ-SF) and Mindful Eating Questionnaire (MEQ-30) were used as data collection tools. There were no group differences between scores of vigorous actives and moderate actives from MEQ-30 total and subsamples in eating control, eating discipline and mindful eating, but there was difference between both groups and inactives group ( $p < 0.05$ ). Inactives group had the highest level in emotional eating; there was group difference between vigorous actives and moderate actives ( $p < 0.05$ ). The results suggest that there are useful items for exploring more about mindful eating, because the differences between mindful eating total and its subsamples and physical activity are important. A better understanding of mindful eating can lead to the development of more effective intervention strategies to address the risk of overweight and obesity in adolescents

**Keywords:** *nutrition, physical activity, mindful eating, obesity.*

## 1. Introduction

The World Health Organization (WHO) has reported that four main factors that lead to cardiovascular deaths are unhealthy eating, physical inactivity, tobacco, alcohol use (WHO, 2014). These unhealthy and changeable behaviors usually occur during adolescence (Croezen et al., 2009). Adolescents, at the age of 15-17, skip breakfast and their physical activity declines (Duarte et al., 2020). For this reason, the period between 15 and 17 should be considered as a critical time to promote healthy lifestyles (Patton et al., 2016) and lifestyle deconstruction and eating habits should be established during this period (Das et al., 2017; Gümüş & Yardımcı, 2020). Nutritional behavior, which is a physiological requirement in childhood, begins to turn into a behavior that can be learned and consciously controlled in adolescence. Sudden increase in burning energy and rapid development play a role in determining the eating habits of adolescents (Uzdilet et al., 2017). Usually, adolescents consume fast food and the number of meals increases during this period. Rapid development leads them to eat frequently and profusely. In addition, food consumption with high sugar and fat content is common nutritional errors in this period (Lassi et al., 2017; Pereira

et al., 2021). These risky behaviors negatively affect the social, emotional, cognitive, and physiological development processes of adolescents (Mairs & Nicholls, 2016). Given that an increase in calorie intake is the basis of obesity, it is extremely important to understand correct factors that influence eating decisions (Nury et al., 2021).

Being overweight or obese in adolescence is associated with lifelong negative health outcomes (Pazin et al., 2020). Excessive consumption of food in adolescence will probably lead to lifelong overweight and obesity (Evensen et al., 2016; Evensen et al., 2017). In addition, obesity in this period is related with the early onset of chronic diseases (Parket et al., 2012) and has negative psychosocial consequences (Wang et al., 2019). Many environmental factors can affect eating, such as hunger, the presence of food, the feeling of eating with others, emotional eating, stressful eating, and boredom (Wansink, 2004; Stroebele & de Castro, 2006; Hemio et al., 2020). Generally, people understand how much they eat and when they will stop eating when they see an empty bowl rather than a feeling of fullness (Guazzelli Williamson et al., 2021). These effects also prevent understanding how much food is eaten (Chen & Antonelli, 2020). Therefore, it is extremely important to develop a better attitude

about the determinants of overweight and obesity in adolescents and to know the health improvement interventions well.

Mindfulness has gained popularity as an adjuvant treatment for adolescents with physical and mental health problems (Lin et al., 2019). Raising awareness for eating behavior is a typical approach applied in behavioral interventions for weight management (Katterman et al., 2014; Stice et al., 2015). The main goal of it is usually to develop acceptance-based behavioral situations, where the focus is on accepting negative experiences and promoting behaviors that are compatible with goals, rather than completely avoiding the negative experiences highlighted in standard behavioral weight management programs (Tronieri et al., 2020). It has been revealed in the studies that energy intake decreases (Chumachenko et al., 2021) and it helps weight control in individuals with improved awareness (Roche et al., 2019). For this reason, it is possible that physical activity creates self-awareness in terms of performance.

Although people accept that environmental factors affect other people, they think that they do not affect them. However, only about 200 unconscious decisions are made per day for food consumption (Wansink, 2004; Moore et al., 2013). If people consciously direct their attention to eating, they may not eat too much. It can prevent them from unhealthiness. In order to understand this, it is vital to determine the state of awareness in which food is consumed. The duration and intensity of physical activity plays an important role precisely at this stage. Because the level of physical activity adjusts food choices with the physiological satiety signaling system (Blundell et al., 2003). In addition, those who do regular physical activity have good nutritional information, and this can change their attitude (Bakhtiar et al., 2021). Nevertheless, those who do regular physical activity like other individuals can get information from the wrong sources (such as the internet; Devlin & Belski, 2015; Trakman et al., 2019). It can lead them to develop improper eating behavior. Therefore, the detection of factors affecting the eating behavior of vigorous active, moderate active or inactive adolescents in the middle adolescence period can be a guide in preventing obesity and obesity-related diseases that they may encounter in their old age. In this study, it was aimed to evaluate whether the duration and intensity of physical activity of adolescents between 15 and 17 changed their nutritional awareness by taking the opinions of their families into consideration.

## 2. Method

### 2.1. Participants and Procedure

In this study, cross-sectional method was used. The dependent variable of the research was the score taken from the scales. Before the data was collected, participants and their families were informed about the purposes and procedures of the research and they signed consent forms stating that they were volunteers. They were told that the study was based on voluntariness, and they could leave the study at any time. It was conducted in accordance with the Declaration of Helsinki and permission was obtained from Usak University Ethics Committee (2021-95 numbered decision). The facts that adolescents are between 15 and 17, they

do not have chronic health problems, mental and physical disabilities that will prevent them from doing physical activity are considered as criteria for participation in the study. 1117 adolescents aged 15–17 years at equal/similar age, education and BMI levels and their parents, living in different cities in Turkey decisively participated in the study. Of the adolescents, 45.1% (504) were female and 54.9% (613) were male. 371 (33.3%) of the respondents were 15, 389 (34.9%) were 16 and 357 (31.8%) were 17. All of the participants were adolescents with a normal body mass index (BMI), as mindful eating and physical activity might have an impact on their results (18.5–24.9). BMI was calculated by dividing weight by height ( $\text{kg/m}^2$ ).

## 2.2. Scales

### 2.2.1. Mindful Eating Questionnaire (MEQ-30)

Mindful Eating Questionnaire (MEQ) was developed by Framson et al. (2009) and it included 28 items and 5 sub factors. It is used to evaluate eating behaviors. It is also practiced in many languages and cultures and a valid and reliable measuring tool (Apolzan et al., 2016; Abdul Basir et al., 2021). MEQ was adopted into Turkish by Köse et al., (2016) and MEQ-30 was developed by adding new questions. Scale has 30 items and 7 subfactors as disinhibition (mindless eating), emotional eating, eating control, mindfulness, eating discipline, conscious nutrition and interference. Questions; 1, 7, 9, 11, 13, 15, 18, 24, 25 and 27 are normally scored. The others are reversely scored \*. \*Reverse scoring is (1=5, 2=4, 3=3, 4=2, 5=1). The items of the questionnaire are scored using a 5-point Likert-style scale as Never (1), Rarely (2), Sometimes (3), Often (4) and Always (5). When scoring the scale, the arithmetic mean of the sub-dimensions and the total score are taken. Score 3 and above taken from the sub-factors indicate that the individual has the characteristics that sub-factor evaluates. The higher score obtained from the scale is, the higher the state of mindful eating is. The Cronbach alpha value of MEQ-30 is 0.733.

### 2.2.2. International Physical Activity Questionnaire Short Form (IPAQ-SF)

IPAQ-SF was developed by Craig et al., (2003) to determine the physical activity levels of everybody in the 15–65. It is applied in many regions of the world (Aibaret et al., 2016; Ekelund et al., 2006; Gallardo et al., 2020). Validity and reliability studies were conducted with the Turkish adaptation of IPAQ-SF (Saglam et al., 2010). When scoring IPAQ-SF, it is based on the fact that the activities are performed at least 10 minutes. After determining the day and minute curves, the MET value was calculated. Physical activity levels after the MET value obtained were classified as vigorous active (healthy-beneficial;  $>3000$  MET min/week), moderate active (600-3000 MET min/week) and inactive ( $<600$  MET-min/week; Ridley et al., 2008; Guidelines for Data Processing and Analysis of the IPAQ-SF, 2005). The answers which contained phrases such as "I don't know," time or days were left incomplete were completely excluded from the analysis. According to the data obtained, 350 people were vigorous active, 398 were moderate active and 359 were inactive. The Cronbach  $\alpha$  internal consistency value of our study is  $\alpha = 0.681$ .

### 2.2.3. Application and Anthropometric Measurements

School administrators, club officials and families who would conduct the study were contacted at the places where measurements and applications would be made before the study. The days and times to go for the measurement were decided in advance. During the specified time period, the adolescent and his/her parent were met at the place where the measurement would be performed. Before the application, the height and weight measurements of adolescents were performed in a separate and privacy-requiring section. Weight measurement was performed with the Jawon Make Body Composition Analyzer (Model IOI-353) with 0.1 accuracy with bare feet when they had light clothing on them. Height measurement was made with a portable stadiometer (Seca Corporation, USA) which had 0.1 cm accuracy. After collecting anthropometric data, participants were taken into the classroom environment in groups (25–30 people). International Physical Activity Questionnaire Short Form (IPAQ-SF) and Mindful Eating Questionnaire (MEQ-30) were projected on the screen respectively. Adolescents and their families were also asked to answer scale questions at the same time. Questions in the IPAQ-SF and MEQ were answered after adolescent and parent agreed. The hand-filled data were collected from the participants for evaluation. There was no section, which include personal information of the participants in IPAQ-SF and MEQ-30 scales.

### 2.3. Statistical Method

SPSS 21.0 was used to analyze the data. Descriptive statistics were given as number, percentage, average and standard deviation. KolmogorowSmirnow test was applied to examine whether the data showed normal distribution. For multiple comparisons in normal distributions One-Way ANOVA Test and for Post- Hoc test Tukey HSD were used. The statistical significance level was considered as  $p < 0.05$ .

### 3. Results

There was no difference between score of vigorous actives and moderate actives from MEQ-30 total and subsamples in eating control, eating discipline and mindful eating, but there was difference between both groups and inactives group ( $p < 0.05$ ). Statistically, no significant difference between disinhibition and mindfulness ( $p > 0.05$ ) was observed, but vigorous active adolescents got the highest scores. The highest level in emotional eating belonged to inactives group, there was difference between vigorous actives and moderate actives ( $p < 0.05$ ). Although there was no difference statistically, inactive adolescents were mostly affected by external factors ( $p > 0.05$ ; Table 1).

**Table 1.** Comparison of Mindful Eating and Sub-Factors According to Physical Activity

	IPAQ-SF	N	M	SD	F	p
Disinhibition	Vigorous active	350	3.12	0.88	3.221	0.059
	Moderate active	398	3.08	0.85		
	Inactive	359	3.08	0.85		
Emotional Eating	Vigorous active	350	3.03 <sup>B</sup>	0.99	7.612	0.001*
	Moderate active	398	3.21 <sup>A</sup>	0.96		
	Inactive	359	3.27 <sup>A</sup>	0.95		
Control of Eating	Vigorous active	350	3.57 <sup>A</sup>	0.82	3.327	0.036*
	Moderate active	398	3.48 <sup>A</sup>	0.84		
	Inactive	359	3.42 <sup>B</sup>	0.74		
Focusing	Vigorous active	350	3.12	0.60	1.413	0.244
	Moderate active	398	3.09	0.54		
	Inactive	359	3.05	0.53		
Eating Discipline	Vigorous active	350	2.91 <sup>A</sup>	0.88	11.230	0.000*
	Moderate active	398	2.88 <sup>A</sup>	0.81		
	Inactive	359	2.66 <sup>B</sup>	0.75		
Mindfulness	Vigorous active	350	3.21 <sup>A</sup>	0.58	2.656	0.040*
	Moderate active	398	3.09 <sup>A</sup>	0.48		
	Inactive	359	3.02 <sup>B</sup>	0.57		
Interference	Vigorous active	350	3.37	1.04	0.464	0.629
	Moderate active	398	3.41	1.07		
	Inactive	359	3.44	0.98		
MEQ-30 Total	Vigorous active	350	3.22 <sup>A</sup>	0.43	4.983	0.007*
	Moderate active	398	3.17 <sup>A</sup>	0.45		
	Inactive	359	3.10 <sup>B</sup>	0.42		

\* $p < 0.05$  means significant difference. A,B: Different letters in the same column indicate the difference between the groups. MEQ-30: Mindful Eating Questionnaire. IPAQ-SF: International Physical Activity Questionnaire-Short Form

## 4. Discussion

Soon after a short period of intense physical activity, a number of disadvantages can be experienced in food choices, leading to delicious and high-energy foods (fatty and sweet; Gustafson et al., 2018; Koehler et al., 2021). Because intense physical activity increases neural responses in reward-related areas of the brain (Schmitt et al., 2020), but it is a temporary case. Physical activity can positively change food consumption due to improved appetite control combined with a higher motivation to engage in healthy behavior (Joo et al., 2019). Cavadini et al. (2000), Croll et al. (2006), and Meng et al. (2018) found that adolescents engaged in physical activity exhibited healthier eating habits than adolescents who did not. There were no group differences between scores of vigorous actives and moderate actives from MEQ-30 total and subsamples in eating control, eating discipline and mindful eating, but there was difference between both groups and inactives group. Statistically, there was no significant difference in mindfulness ( $p>0.05$ ) factor, but vigorous active adolescents got the highest scores. These results show that physical activity during adolescence has a significant effect on eating. Appetite control develops and positive changes in food preferences occur in accordance with an overall improvement in parallel with physical activity training (Beaulieu et al., 2020). Knowing the effect of conscious eating on sports performance motivates those who do physical activity for healthy eating (Stokes et al., 2018). Friends, coaches, dieticians in the sports community have a positive effect on healthy eating and they can be factors that positively affect eating control, eating discipline, and mindful eating (Eckert et al., 2021; Hackman et al., 1992; Kalavana et al., 2010). These resources can enable adolescent athletes to exhibit healthy eating behavior.

Emotional eating behavior (eating in response to negative emotions) usually occurs during adolescence and can lead to excessive energy consumption (sugar; van Strien, 2018). Emotional eating is either a learned behavior affected by genetic factors compared to environmental factors (Russell, & Russell, 2020) or it can occur when the mechanisms that control food intake stimulates the sensory pleasures of eating delicious food (Finlayson et al., 2009). The results are consistent with the research showing the links between emotional eating deceptions and stress, poor food selection, and diet quality (Heo et al., 2018; Wallace et al., 2021). According to the results of our study, inactives group had the highest level in emotional eating; there was difference between vigorous actives and moderate actives ( $p<0.05$ ). Although there was no difference statistically, inactive adolescents were mostly affected by external factors. According to these results, physical activity is one of the most important factors to be considered in terms of enabling eating regulation. In other words, psychological stress can have a significant regulatory effect on eating and emotional functioning in addition to reducing lubrication through energy balance (Messerli-Bürgy et al., 2019; Smith et al., 2021). It is confirmed with the outcomes of the present study that the more physical activity is related with the more mental health (more positive effect), better appetite regulation, less food sensation, less binge eating (Cadenas-Sanchez et al., 2021; Luo et al., 2018). Increased positive effect and decreased negative effect due to physical activity may have

revealed impacts on eating and appetite (Liao et al., 2015), because physical activity can affect body regulation at both individual and instant levels through its effects on emotional eating behavior (Smith et al., 2021). Physical activity can regulate eating behavior by reducing food-craving activity in brain reward regions (Blanchet et al., 2018; Luo et al., 2018). Increased physical activity can achieve this by suppressing the hedonic urge on overeating (Joseph et al., 2011).

Interpreting the results, it should be taken into account that there are some strengths and limitations in this study. The study sample represented a large population that was fairly well balanced (age, gender, education). Although it consisted largely equal male and female participants, the balance within the groups was not clear. Providing more balanced groups in terms of gender could contribute more to the results of the study. Our study is cross-sectional and inferences about causal relationships cannot be made. On the other hand, having two different variables (adolescents and their families) strengthens the accuracy of the data, especially adding validity to reports of adolescent awareness and eating habits reported by parents.

While our study results indicate the impact of physical activity on eating control, eating discipline and mindful eating among MEQ-30 total and sub factors, there is no statistical difference and PA may have a positive effect on emotional eating activity and negative eating behavior towards external factors. The results suggest that there are useful outcomes for exploring more about eating awareness. A better understanding of mindful eating can lead to the development of more effective intervention strategies to address the risk of overweight and obesity in adolescents.

### Acknowledgements

The author would like to thank all the participants in the study.

### Disclosure Statement

The authors declared that there is no potential conflict of interest.

### Funding

No funding to declare.

## References

- Abdul Basir, S. M., Abdul Manaf, Z., Ahmad, M., Abdul Kadir, N. B., Ismail, W., Mat Ludin, A. F., & Shahr, S. (2021). Reliability and validity of the Malay Mindful Eating Questionnaire (MEQ-M) among overweight and obese adults. *International Journal of Environmental Research and Public Health*, 18(3), 1021.
- Aibar, A., García-González, L., Abarca-Sos, A., Murillo, B., & Zaragoza, J. (2016). Analizando la validación del International Physical Activity Questionnaire en jóvenes adolescentes: Un protocolo modificado para la recogida de los datos. *SPORT TK-Revista EuroAmericana de Ciencias del Deporte*, 5(2), 123–132.
- Apolzan, J. W., Myers, C. A., Cowley, A. D., Brady, H., Hsia, D. S., Stewart, T. M., Redman, L. M., & Martin, C. K. (2016). Examination of the reliability and validity of the Mindful Eating Questionnaire in pregnant women. *Appetite*, 100, 142–151.
- Bakhtiar, M., Masud-Ur-Rahman, M., Kamruzzaman, M., Sultana, N., & Rahman, S. S. (2021). Determinants of nutrition knowledge, attitude and practices of adolescent sports trainees: A cross-sectional study in Bangladesh. *Heliyon*, 7(4), e06637.
- Beaulieu, K., Oustric, P., & Finlayson, G. (2020). The impact of physical activity on food reward: Review and conceptual synthesis of evidence

- from observational, acute, and chronic exercise training studies. *Current Obesity Reports*, 9(2), 63–80.
- Blanchet, C., Mathieu, M. È., St-Laurent, A., Fecteau, S., St-Amour, N., & Drapeau, V. (2018). A systematic review of physical activity interventions in individuals with binge eating disorders. *Current Obesity Reports*, 7(1), 76–88.
- Cadenas-Sanchez, C., Mena-Molina, A., Torres-Lopez, L. V., Migueles, J. H., Rodriguez-Ayllon, M., Lubans, D. R., & Ortega, F. B. (2021). Healthier minds in fitter bodies: A systematic review and meta-analysis of the association between physical fitness and mental health in youth. *Sports Medicine*, 51(12), 2571–2605. 10.1007/s40279-021-01520-y
- Cavadini, C., Decarli, B., Grin, J., Narring, F., & Michaud, P. A. (2000). Food habits and sport activity during adolescence: Differences between athletic and non-athletic teenagers in Switzerland. *European Journal of Clinical Nutrition*, 54(1), S16–S20.
- Chen, P. J., & Antonelli, M. (2020). Conceptual models of food choice: Influential factors related to foods, individual differences, and society. *Foods*, 9(12), 1898.
- Chumachenko, S. Y., Cali, R. J., Rosal, M. C., Allison, J. J., Person, S. J., Ziedonis, D., Nephew, B. C., Moore, C. M., Zhang, N., King, J. A., & Fulwiler, C. (2021). Keeping weight off: Mindfulness-based stress reduction alters amygdala functional connectivity during weight loss maintenance in a randomized control trial. *PLoS one*, 16(1), e0244847.
- Craig, C. L., Marshall, A. L., Sjoström, M., Bauman, A. E., Booth, M. L., Ainsworth, B. E., Pratt, M., Ekelund, U., Yngve, A., Sallis, J. F., Oja, P. (2003) International physical activity questionnaire: 12-country reliability and validity. *Medicine and Science in Sports and Exercise*, 35(8), 1381–1395.
- Croezen, S., Visscher, T. L., Ter Bogt, N. C., Veling, M. L., & Haveman-Nies, A. (2009). Skipping breakfast, alcohol consumption and physical inactivity as risk factors for overweight and obesity in adolescents: results of the E-MOVO project. *European Journal of Clinical Nutrition*, 63(3), 405–412.
- Croll, J. K., Neumark-Sztainer, D., Story, M., Wall, M., Perry, C., & Harnack, L. (2006). Adolescents involved in weight-related and power team sports have better eating patterns and nutrient intakes than non-sport-involved adolescents. *Journal of the American Dietetic Association*, 106(5), 709–717.
- Das, J. K., Salam, R. A., Thornburg, K. L., Prentice, A. M., Campisi, S., Lassi, Z. S., Koletzko, B., Bhutta, Z., A. (2017). Nutrition in adolescents: Physiology, metabolism, and nutritional needs. *Annals of the New York Academy of Sciences*, 1393(1), 21–33.
- Devlin, B. L., & Belski, R. (2015). Exploring general and sports nutrition and food knowledge in elite male Australian athletes. *International Journal of Sport Nutrition and Exercise Metabolism*, 25(3), 225–232.
- Doro Altan, A. M., MancaBitti, M. L., Buonomo, E., Scarcella, P., Mancinelli, S., Arcano, S., & Palombi, L. (2008). Un intervento di sanità pubblica per la valutazione e la prevenzione del diabete di tipo 2 e delle malattie cardiovascolari in bambini e adolescenti obesi [Evaluation and prevention of type II diabetes mellitus and cardiovascular diseases in obese children and adolescents: a public health intervention in a local health organisation in Rome (Italy)]. *Igiene e Sanità Pubblica*, 64(3), 345–360.
- Duarte, L., Fujimori, E., Borges, A. L., Kurihayashi, A., Steen, M., & Roman Lay, A. (2020). Body weight dissatisfaction is associated with cardiovascular health-risk behaviors among Brazilian adolescents: Findings from a national survey. *International Journal of Environmental Research and Public Health*, 17(23), 8929.
- Eck, K. M., & Byrd-Bredbenner, C. (2021). Food choice decisions of collegiate Division I athletes: A qualitative exploratory study. *Nutrients*, 13(7), 2322.
- Ekelund, U., Sepp, H., Brage, S., Becker, W., Jakes, R., Hennings, M., & Wareham, N. J. (2006). Criterion-related validity of the last 7-day, short form of the International Physical Activity Questionnaire in Swedish adults. *Public Health Nutrition*, 9(2), 258–265.
- Evensen, E., Emaus, N., Kokkvol, A., Wilsgaard, T., Furberg, A. S., & Skeie, G. (2017). The relation between birthweight, childhood body mass index, and overweight and obesity in late adolescence: a longitudinal cohort study from Norway, The Tromsø Study, Fit Futures. *BMJ open*, 7(6), e015576.
- Evensen, E., Wilsgaard, T., Furberg, A. S., & Skeie, G. (2016). Tracking of overweight and obesity from early childhood to adolescence in a population-based cohort - the Tromsø Study, Fit Futures. *BMC Pediatrics*, 16, 64.
- Finlayson, G., Bryant, E., Blundell, J. E., & King, N. A. (2009). Acute compensatory eating following exercise is associated with implicit hedonic wanting for food. *Physiology & Behavior*, 97(1), 62–67.
- Framson, C., Kristal, A. R., Schenk, J. M., Littman, A. J., Zeliadt, S., & Benitez, D. (2009). Development and validation of the mindful eating questionnaire. *Journal of the American Dietetic Association*, 109(8), 1439–1444.
- Gallardo, L. O., Abarca-Sos, A., & Doña, A. M. (2020). Expectancy-value model related to physical activity behaviors in Chilean and Spanish adolescents. *International Journal of Environmental Research and Public Health*, 17(21), 8219.
- Guazzelli Williamson, V., Lee, A. M., Miller, D., Huo, T., Maner, J. K., & Cardel, M. (2021). Psychological resilience, experimentally manipulated social status, and dietary intake among adolescents. *Nutrients*, 13(3), 806.
- Gustafson, C. R., Rakhmatullaeva, N., Beckford, S. E., Ammachathram, A., Cristobal, A., & Koehler, K. (2018). Exercise and the timing of snack choice: Healthy snack choice is reduced in the post-exercise state. *Nutrients*, 10(12), 1941.
- Gümüş, A. B., Yardımcı, H. (2020). Erken ve orta adölesan dönemdeki bireylerin ana öğün tüketim durumları ve yeme tutumları. *Ankara Sağlık Bilimleri Dergisi (ASBD)*, 9(1), 176–184.
- Hackman, R. M., Katra, J. E., & Geertsens, S. M. (1992). The athletic trainer's role in modifying nutritional behaviors of adolescent athletes: Putting theory into practice. *Journal of Athletic Training*, 27(3), 262–267.
- Hemio, K., Lindström, J., Peltonen, M., Härmä, M., Viitasalo, K., & Puttonen, S. (2020). The association of work stress and night work with nutrient intake - a prospective cohort study. *Scandinavian Journal of Work, Environment & Health*, 46(5), 533–541.
- Heo, M., Jimenez, C. C., Lim, J., Isasi, C. R., Blank, A. E., Lounsbury, D. W., Fredericks, L., Bouchard, M., Faith, M. S., & Wylie-Rosett, J. (2018). Effective nationwide school-based participatory extramural program on adolescent body mass index, health knowledge and behaviors. *BMC Pediatrics*, 18(1), 7.
- International Physical Activity Questionnaire. (2005). Guidelines for Data Processing and Analysis of the International Physical Activity Questionnaire (IPAQ)-Short and Long Forms. <http://www.ipaq.ki.se/scoring.pdf>
- Joo, J., Williamson, S. A., Vazquez, A. I., Fernandez, J. R., & Bray, M. S. (2019). The influence of 15-week exercise training on dietary patterns among young adults. *International Journal of Obesity*, 43(9), 1681–1690.
- Joseph, R. J., Alonso-Alonso, M., Bond, D. S., Pascual-Leone, A., & Blackburn, G. L. (2011). The neurocognitive connection between physical activity and eating behaviour. *Obesity Reviews*, 12(10), 800–812.
- Kalavana, T. V., Maes, S., & De Gucht, V. (2010). Interpersonal and self-regulation determinants of healthy and unhealthy eating behavior in adolescents. *Journal of Health Psychology*, 15(1), 44–52.
- Katterman, S. N., Kleinman, B. M., Hood, M. M., Nackers, L. M., & Corsica, J. A. (2014). Mindfulness meditation as an intervention for binge eating, emotional eating, and weight loss: A systematic review. *Eating Behaviors*, 15(2), 197–204.
- Koehler, K., Beckford, S. E., Thayer, E., Martin, A. R., Boron, J. B., & Stevens, J. R. (2021). Exercise shifts hypothetical food choices toward greater amounts and more immediate consumption. *Nutrients*, 13(2), 347.
- Köse, G., Tayfur, M., Birincioglu, I., Dönmez, A. (2016). Adaptation study of the Mindful Eating Questionnaire (MEQ) into Turkish. *Journal of Cognitive Behavioral Psychotherapy and Research*, 5(3), 125–134.
- Lassi, Z. S., Mansoor, T., Salam, R. A., Bhutta, S. Z., Das, J. K., Bhutta, Z. A. (2017). Review of nutrition guidelines relevant for adolescents in low- and middle-income countries. *Annals of the New York Academy of Sciences*, 1393(1), 51–60.
- Liao, Y., Shonkoff, E. T., & Dunton, G. F. (2015). The acute relationships between affect, physical feeling states, and physical activity in daily life: A review of current evidence. *Frontiers in Psychology*, 6, 1975.

- Luo, S., O'Connor, S. G., Belcher, B. R., & Page, K. A. (2018). Effects of physical activity and sedentary behavior on brain response to high-calorie food cues in young adults. *Obesity*, 26(3), 540–546.
- Mairs, R., & Nicholls, D. (2016). Assessment and treatment of eating disorders in children and adolescents. *Archives of Disease in Childhood*, 101(12), 1168–1175.
- Meng, Y., Manore, M. M., Schuna, J. M., Jr, Patton-Lopez, M. M., Branscum, A., & Wong, S. S. (2018). Promoting healthy diet, physical activity, and life-skills in high school athletes: Results from the WAVE ripples for change childhood obesity prevention two-year intervention. *Nutrients*, 10(7), 947.
- Messerli-Bürge, N., Horsch, A., Schindler, C., Boichat, A., Kriemler, S., Munsch, S., Crottet, B., Marquez-Vidal, P. M., Borghini, A., & Puder, J. J. (2019). Influence of acute physical activity on stress reactivity in obese and normal weight children: A randomized controlled trial. *Obesity Facts*, 12(1), 115–130.
- Monroe J. T. (2015). Mindful eating: Principles and practice. *American Journal of Lifestyle Medicine*, 9(3), 217–220.
- Moor, K. R., Scott, A. J., & McIntosh, W. D. (2013). Mindful eating and its relationship to body mass index and physical activity among university students. *Mindfulness*, 4(3), 269–274.
- Nury, E., Morze, J., Grummich, K., Rücker, G., Hoffmann, G., Angele, C. M., Steinacker, J. M., Conrad, J., Schmid, D., Meerpohl, J. J., & Schwingshackl, L. (2021). Effects of nutrition intervention strategies in the primary prevention of overweight and obesity in school settings: A protocol for a systematic review and network meta-analysis. *Systematic Reviews*, 10(1), 122.
- Park, M. H., Falconer, C., Viner, R. M., & Kinra, S. (2012). The impact of childhood obesity on morbidity and mortality in adulthood: A systematic review. *Obesity Reviews*, 13(11), 985–1000.
- Patton, G. C., Sawyer, S. M., Santelli, J. S., Ross, D. A., Afifi, R., Allen, N. B., Arora, M., Azzopardi, P., Baldwin, W., Bonell, C., Kakuma, R., Kennedy, E., Mahon, J., McGovern, T., Mokdad, A. H., Patel, V., Petroni, S., Reavley, N., Taiwo, K., Waldfogel, J., ... Viner, R. M. (2016). Our future: Alancet commission on adolescent health and wellbeing. *Lancet*, 387(10036), 2423–2478.
- Pazin, D. C., da Luz Kaestner, T. L., Olandoski, M., Baena, C. P., de Azevedo Abreu, G., Kuschner, M., Bloch, K. V., & Faria-Neto, J. R. (2020). Association between abdominal waist circumference and blood pressure in Brazilian adolescents with normal body mass index: Waist circumference and blood pressure in Adolescents. *Global Heart*, 15(1), 27.
- Pereira, J. L., de Castro, M. A., Crispim, S. P., Fisberg, R. M., Isasi, C. R., Mossavar-Rahmani, Y., Van Horn, L., Carnethon, M. R., Daviglus, M. L., Perreira, K. M., Gallo, L. C., Sotres-Alvarez, D., & Mattei, J. (2021). Comparing methods from the national cancer institute vs multiple source method for estimating usual intake of nutrients in the Hispanic community health study/study of Latino youth. *Journal of the Academy of Nutrition and Dietetics*, 121(1), 59–73.e16.
- Ridley, K., Ainsworth, B. E., & Olds, T. S. (2008). Development of a compendium of energy expenditures for youth. *The International Journal of Behavioral Nutrition and Physical Activity*, 5, 45.
- Roche, A. I., Kroska, E. B., & Denburg, N. L. (2019). Acceptance- and mindfulness-based interventions for health behavior change: Systematic reviews and meta-analyses. *Journal of Contextual Behavioral Science*, 13, 74–93.
- Russell, C. G., & Russell, A. (2020). "Food" and "non-food" self-regulation in childhood: A review and reciprocal analysis. *The International Journal of Behavioral Nutrition and Physical Activity*, 17(1), 33.
- Sacks, G., & Looi, E. (2020). The advertising policies of major social media platforms overlook the imperative to restrict the exposure of children and adolescents to the promotion of unhealthy foods and beverages. *International Journal of Environmental Research and Public Health*, 17(11), 4172.
- Saglam, M., Arikan, H., Savci, S., Inal-Ince, D., Bosnak-Guclu, M., Karabulut, E., & Tokgozoglu, L. (2010). International physical activity questionnaire: Reliability and validity of the Turkish version. *Perceptual and Motor Skills*, 111(1), 278–284.
- Schmitt, A., Upadhyay, N., Martin, J. A., Rojas Vega, S., Strüder, H. K., & Boecker, H. (2020). Affective modulation after high-intensity exercise is associated with prolonged amygdala-insular functional connectivity increase. *Neural Plasticity*, 2020, 7905387.
- Schneider D. (2000). International trends in adolescent nutrition. *Social Science & Medicine*, 51(6), 955–967.
- Smith, K. E., O'Connor, S. M., Mason, T. B., Wang, S., Dzibur, E., Crosby, R. D., Wonderlich, S. A., Salvy, S. J., Feda, D. M., & Roemmich, J. N. (2021). Associations between objective physical activity and emotional eating among adiposity-discordant siblings using ecological momentary assessment and accelerometers. *Pediatric Obesity*, 16(3), e12720.
- Stice, E., Rohde, P., Butryn, M., Menke, K. S., & Marti, C. N. (2015). Randomized controlled pilot trial of a novel dissonance-based group treatment for eating disorders. *Behaviour Research and Therapy*, 65, 67–75.
- Stokes, E. G., Hughes, R., Shaw, D. M., O'Connor, H. T., & Beck, K. L. (2018). Perceptions and determinants of eating for health and performance in high-level male adolescent rugby union players. *Sports*, 6(2), 49.
- Stroebele, N., & de Castro, J. M. (2006). Influence of physiological and subjective arousal on food intake in humans. *Nutrition*, 22(10), 996–1004.
- Trakman, G. L., Forsyth, A., Hoyer, R., & Belski, R. (2019). Australian team sports athletes prefer dietitians, the internet and nutritionists for sports nutrition information. *Nutrition & Dietetics*, 76(4), 428–437.
- Tronieri, J. S., Wadden, T. A., Pearl, R. L., Berkowitz, R. I., Alamuddin, N., & Chao, A. M. (2020). Mindful eating, general mindful awareness, and acceptance as predictors of weight loss. *Mindfulness*, 11(12), 2818–2827.
- Uzdil, Z., Özenoğlu, A., Ünal, G. (2017). Lise öğrencilerinde yeme tutumlarının beslenme alışkanlıkları, antropometrik ve demografik özellikleri ile ilişkisi. *Düzce Üniversitesi Sağlık Bilimleri Enstitüsü Dergisi*, 7(1), 11–18.
- van Strien T. (2018). Causes of emotional eating and matched treatment of obesity. *Current Diabetes Reports*, 18(6), 35.
- Wallace, D. D., Lytle, L. A., Albrecht, S., & Barrington, C. (2021). All of that causes me stress: An exploration of the sources of stress experienced by Latinxs living with prediabetes. *Journal of Latina/o Psychology*, 9(3), 204–216.
- Wang, S., Sun, Q., Zhai, L., Bai, Y., Wei, W., & Jia, L. (2019). The prevalence of depression and anxiety symptoms among overweight/obese and non-overweight/non-obese children/adolescents in China: A systematic review and meta-analysis. *International Journal of Environmental Research and Public Health*, 16(3), 340.
- Wansink B. (2004). Environmental factors that increase the food intake and consumption volume of unknowing consumers. *Annual review of nutrition*, 24, 455–479.
- World Health Organization (2014). *Global status report on noncommunicable diseases 2014*. World Health Organization. <https://apps.who.int/iris/handle/10665/148114>

#### Suggested Citation:

AKIL, M. (2022). The mediating role of duration and intensity of physical activity in increasing eating awareness: Adolescent and parental participation study, *Journal of Teaching, Research, and Media in Kinesiology*, 8, 10–15.